



Organic Chemistry 1

Dr Muhannad Amer

Summer

2017

Chapter 1 Bonding and Isomerism

Organic Chemistry is the chemistry of carbon compounds

Why does sucrose melt at 185°C while table salt melts at 801°C ?

Why do both substances dissolve in water and olive oil does not?

Why does methyl butyrate smell like pears while propyl acetate smell like apple yet they have the same number and kind of atoms?


Bonding is the key to the structure, physical properties and chemical behavior of different kinds of matter.

1.1 How Electrons are arranged in Atom

- The Structure of an Atom
- An atom consists of electrons, positively charged protons, and neutral neutrons
- Electrons form chemical bonds
- Atomic number: numbers of protons in its nucleus
- Mass number: the sum of the protons and neutrons of an atom
- Isotopes have the same atomic number but different mass numbers
- The atomic weight: the average weighted mass of its atoms
- Molecular weight: the sum of the atomic weights of all the atoms in the molecule



Mass Number and Isotope



● Proton ● Neutron ● Electron

Carbon atoms

Mass Number → $^{12}_6\text{C}$, $^{13}_6\text{C}$
Atomic Number → ^6_6C , ^6_6C

Atomic Number = The number of protons
= (The number of electrons)

Mass Number = The number of protons + The number of neutrons

Carbon has two isotopes: ^{12}C and ^{13}C . Both show the same chemical properties because they have the same atomic number 6. But the mass of ^{13}C is larger than that of ^{12}C , because ^{13}C has one more neutron.

The ground-state electronic configuration describes the orbitals occupied by the atom's electrons with the lowest energy

Table 1.2  Electron Arrangements of the First 18 Elements

Atomic number	Element	Number of electrons in each orbital				
		1s	2s	2p	3s	3p
1	H	1				
2	He	2				
3	Li	2	1			
4	Be	2	2			
5	B	2	2	1		
6	C	2	2	2		
7	N	2	2	3		
8	O	2	2	4		
9	F	2	2	5		
10	Ne	2	2	6		
11	Na	2	2	6	1	
12	Mg	2	2	6	2	
13	Al	2	2	6	2	1
14	Si	2	2	6	2	2
15	P	2	2	6	2	3
16	S	2	2	6	2	4
17	Cl	2	2	6	2	5
18	Ar	2	2	6	2	6

Table 1.3 Valence Electrons of the First 18 Elements


Group	I	II	III	IV	V	VI	VII	VIII
	H·							He:
	Li·	·Be·	·B·	·C·	·N:	·O:	:F:	:Ne:
	Na·	·Mg·	·Al·	·Si·	·P:	·S:	:Cl:	:Ar:


1.2 Ionic and Covalent bonding

Lewis's theory: an atom will give up, accept, or share electrons in order to achieve a filled outer shell or an outer shell that contains eight electrons

Ionic Compounds

are composed of positively charged cations and negatively charged anions





Sodium chloride, NaCl, is an ionic crystal. The purple spheres represent sodium ions, Na, and the green spheres are chloride ions, Cl₂.

Each ion is surrounded by six oppositely charged ions, except for those ions that are at the surface of the crystal.

The Covalent Bond

Covalent bonds are formed by sharing electrons



Bond energy (BE) is the energy necessary to break a mole of covalent bonds. The amount of energy depends on the type of bond broken.

The bond length is the average distance between two covalently bonded atoms.


Problem 1.4

Write an equation for the formation of chlorine molecule




1.4 Carbon-Carbon Single Bonds


Carbon could share electrons with not only different elements but also carbon.




or



ethane




or



hexachloroethane

Less heat is required to break the C-C bond in ethane than the H-H bond in a hydrogen molecule. The C-C-bond in ethane is 1.54 Å. The H-H bond in H₂ molecule is 0.74 Å. The C-H is about 1.09 Å, close to the average of H-H bond and C-C bond.



A radical is a molecular fragment with an odd number of electrons

1.4 Polar Covalent Bonds

Is a covalent bond in which the electron are is not shared equally between the atoms

The bond polarization is indicated by an arrow whose head is negative and whose tail is marked with a plus sign. Alternatively, a partial charge, written as $\delta+$ or $\delta-$.





Table 1.4  **Electronegativities of Some Common Elements**

Group

I	II	III	IV	V	VI	VII
H 2.2						
Li 1.0	Be 1.6	B 2.0	C 2.5	N 3.0	O 3.4	F 4.0
Na 0.9	Mg 1.3	Al 1.6	Si 1.9	P 2.2	S 2.6	Cl 3.2
K 0.8	Ca 1.0					Br 3.0
						I 2.7


 < 1.0

 1.0–1.4


 1.5–1.9

 2.0–2.4

 2.5–2.9

 3.0–3.4

Bond polarization in tetrachloromethane



Problem 1.10


Predict the polarity of the P-Cl bond and the S-O bond

1.6 Multiple Covalent Bonds



A

or






B

or



C



or



or



hydrogen cyanide

1.7 Valance


The valence of an element is simply the number of bonds that an atom of the element can form. The number is normally equal to the number of electron needed to fill the valence shell.

Table 1.5 ▀ Valences of Common Elements

Element	H·	· $\overset{\cdot}{\underset{\cdot}{\text{C}}}$ ·	· $\overset{\cdot}{\underset{\cdot}{\text{N}}}$:	· $\overset{\cdot\cdot}{\underset{\cdot}{\text{O}}}$:	: $\overset{\cdot\cdot}{\underset{\cdot}{\text{F}}}$:	: $\overset{\cdot\cdot}{\underset{\cdot}{\text{Cl}}}$:
Valence	1	4	3	2	1	1

1.8 Isomerism

Isomers are molecules with the same molecular formula but different arrangement of atoms




ethanol

(ethyl alcohol)

bp 78.5°C

and



methoxymethane

(dimethyl ether)



bp -23.6°C

Structural (or constitutional) isomers






are the compounds that have the same molecular formula but different structural formulas.

Problem 1.20

Draw structural formulas for the three possible isomers of C_3H_8O






Suppose we keep the chain of four carbons and try to connect the fifth carbon somewhere else.





2,2-dimethylpropane, bp 10°C
(neopentane)

PROBLEM 1.21

To which isomer of C_5H_{12} does each of the following structural formulas correspond?



1.10 Abbreviated Structural Formula






n-pentane



isopentane



neopentane



n-pentane




isopentane




neopentane



n-pentane




isopentane





neopentane

Each line segment have a carbon atom at each end




Example 1.12

Write a more detailed structural formula for



or




1.11 Formal Charge

The formal charge on an atom in a covalently bonded molecule or ion is the number of valence electrons in the neutral atom minus the number of covalent bonds to the atom and the number of unshared electrons on the atom.

$$\text{Formal charge} = \frac{\text{number of valence electrons in the neutral atom}}{\text{number of valence electrons in the neutral atom}} - \left(\text{unshared electrons} + \frac{\text{half the shared electrons}}{\text{electrons}} \right)$$

or, in a simplified form,

$$\text{Formal charge} = \frac{\text{number of valence electrons in the neutral atom}}{\text{number of valence electrons in the neutral atom}} - (\text{dots} + \text{bonds})$$



hydronium ion

For H atom

$$\text{Formal charge} = 1 - (0 + 1) = 0$$


For O atom

$$\text{Formal charge} = 6 - (2 + 3) = 1 \quad +1$$

Problem 1.25


Calculate the formal charge on the nitrogen atom in ammonia, NH_3 ; in the ammonium ion, NH_4^+ ; and in the amide ion, NH_2^-

ammonia




$$\text{formal charge on nitrogen} = 5 - (2 + 3) = 0$$

ammonium ion



$$\text{formal charge on nitrogen} = 5 - (0 + 4) = +1$$

amide ion





$$\text{formal charge on nitrogen} = 5 - (4 + 2) = -1$$

The formal charge on hydrogen in all three cases is zero [$1 - (0 + 1) = 0$].


1.12 Resonance

Sometimes, an electron pair is involved with more than two atoms. Molecules and ions in which this occurs can not be adequately represented by a single electron-dot structure. Please consider the structure of the carbonate ion, CO_3^{2-} .



- Only electrons can be moved (usually lone pairs or pi electrons).
- All the bond lengths are the same.
- The real structure is a resonance hybrid.

- Physical measurement tell us that all three C-O bond length are identical: 1.31 Angstrom (Å).
- This distance is between the normal C=O (1.20 Å) and C-O (1.41 Å).
- We usually say the real carbonate ion has a structure that is resonance hybrid of the three contributing resonance structures.




carbonate ion
resonance hybrid


1.13 Arrow Formalism


Arrow system is very important in Chemistry and has specific meaning.





Curved arrows  a pair of electron moving

Fishhook arrows  single electron moving

Straight arrows  point from reactants to products in chemical reaction equations


Straight arrow with half-heads  used in pairs to indicate that the reaction is reversible.

double-headed straight arrow  between two structures indicates that they are resonance structure




The Orbital View of Bonding; the Sigma Bond

The shapes of the *s* and *p* orbitals used by the valence electrons of carbon. The nucleus is at the origin of the three coordinate axes.





The molecular orbital representation of covalent bond formation between two hydrogen atoms.




Such orbitals are called sigma (σ) orbitals, and the bond is referred to as a sigma bond. Sigma bonds may also be formed by the overlap of an s and a p orbital or of two p orbitals,



Carbon sp^3 Hybrid Orbitals




sp^3



Tetrahedral Carbon; the Bonding in Methane

A molecule of methane, CH_4 , is formed by the overlap of the four sp^3 carbon orbitals with the $1s$ orbitals of four hydrogen atoms. The resulting molecule has the geometry of a regular tetrahedron and contains four sigma bonds of the sp^3-s type.



1.18 Classification According to Functional Group

Table 1.6 The Main Functional Groups








	Structure	Class of compound	Specific example	Common name of the specific example
<i>A. Functional groups that are a part of the molecular framework</i>		alkane	CH ₃ —CH ₃	ethane, a component of natural gas
		alkene	CH ₂ =CH ₂	ethylene, used to make polyethylene
		alkyne	HC≡CH	acetylene, used in welding
		arene		benzene, raw material for polystyrene and phenol
<i>B. Functional groups containing oxygen</i> <i>1. With carbon–oxygen single bonds</i>		alcohol	CH ₃ CH ₂ OH	ethyl alcohol, found in beer, wines, and liquors
		ether	CH ₃ CH ₂ OCH ₂ CH ₃	diethyl ether, once a common anesthetic

Table 1.6 ■ continued

	Structure	Class of compound	Specific example	Common name of the specific example
2. With carbon–oxygen double bonds*	$\begin{array}{c} \text{O} \\ \\ -\text{C}-\text{H} \end{array}$	aldehyde	$\text{CH}_2=\text{O}$	formaldehyde, used to preserve biological specimens
	$\begin{array}{c} \text{O} \\ \\ -\text{C}-\text{C}-\text{C}- \\ \quad \quad \end{array}$	ketone	$\begin{array}{c} \text{O} \\ \\ \text{CH}_3\text{CCH}_3 \end{array}$	acetone, a solvent for varnish and rubber cement
3. With single and double carbon–oxygen bonds	$\begin{array}{c} \text{O} \\ \\ -\text{C}-\text{OH} \end{array}$	carboxylic acid	$\begin{array}{c} \text{O} \\ \\ \text{CH}_3\text{C}-\text{OH} \end{array}$	acetic acid, a component of vinegar
	$\begin{array}{c} \text{O} \\ \\ -\text{C}-\text{O}-\text{C}- \\ \quad \end{array}$	ester	$\begin{array}{c} \text{O} \\ \\ \text{CH}_3\text{C}-\text{OCH}_2\text{CH}_3 \end{array}$	ethyl acetate, a solvent for nail polish and model airplane glue

<i>C. Functional groups containing nitrogen**</i>	$\begin{array}{c} \\ -\text{C}-\text{NH}_2 \\ \end{array}$	primary amine	$\text{CH}_3\text{CH}_2\text{NH}_2$	ethylamine, smells like ammonia
	$-\text{C}\equiv\text{N}$	nitrile	$\text{CH}_2=\text{CH}-\text{C}\equiv\text{N}$	acrylonitrile, raw material for making Orlon
<i>D. Functional group with oxygen and nitrogen</i>	$\begin{array}{c} \text{O} \\ \\ -\text{C}-\text{NH}_2 \end{array}$	primary amide	$\begin{array}{c} \text{O} \\ \\ \text{H}-\text{C}-\text{NH}_2 \end{array}$	formamide, a softener for paper
<i>E. Functional group with halogen</i>	$-\text{X}$	alkyl or aryl halide	CH_3Cl	methyl chloride, refrigerant and local anesthetic
<i>F. Functional groups containing sulfur†</i>	$\begin{array}{c} \\ -\text{C}-\text{SH} \\ \end{array}$	thiol (also called mercaptan)	CH_3SH	methanethiol, has the odor of rotten cabbage
	$\begin{array}{c} \quad \\ -\text{C}-\text{S}-\text{C}- \\ \quad \end{array}$	thioether (also called sulfide)	$(\text{CH}_2=\text{CHCH}_2)_2\text{S}$	diallyl sulfide, has the odor of garlic

HOMEWORK 1

31- 43

45- 51